

DISCOVERY OF CARBON RADIO RECOMBINATION LINES IN M82

LEAH K. MORABITO¹, J. B. R. OONK^{1,2}, FRANCISCO SALGADO¹, M. CARMEN TORIBIO², H. J. A. RÖTTGERING¹,
A. G. G. M. TIELENS¹, RAINER BECK³, BJÖRN ADEBAHR³, PHILIP BEST⁴, ROBERT BESWICK⁵, ANNALISA BONAFEDE⁶,
GIANFRANCO BRUNETTI⁷, MARCUS BRÜGGEN⁶, KRZYSZTOF T. CHYŻY⁸, J. E. CONWAY⁹, WIM VAN DRIEL¹⁰,
JONATHAN GREGSON¹¹, MARIJKE HAVERKORN^{1,12}, GEORGE HEALD^{2,13}, CATHY HORELLOU⁹, ANDREAS HORNEFFER³,
MARCO IACOBELLI², MATT J. JARVIS^{14,15}, IVAN MARTI-VIDAL⁹, GEORGE MILEY¹, D. D. MULCAHY¹⁶,
EMANUELA ORRÚ^{2,12}, ROBERTO PIZZO², A. M. M. SCAIFE¹⁶, ESKIL VARENIUS⁹, REINOUT J. VAN WEEREN¹⁷,
GLENN J. WHITE^{11,18}, AND MICHAEL W. WISE²

¹ Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, The Netherlands; morabito@strw.leidenuniv.nl

² Netherlands Institute for Radio Astronomy (ASTRON), Postbus 2, 7990 AA Dwingeloo, The Netherlands

³ MPI für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

⁴ SUPA, Institute for Astronomy, Royal Observatory Edinburgh, Blackford Hill, Edinburgh EH9 3HJ, UK

⁵ Jodrell Bank Centre for Astrophysics, University of Manchester, Oxford Road, Manchester M13 9PL, UK

⁶ Hamburg Observatory, University of Hamburg, Gojenbergsweg 112, D-21029 Hamburg, Germany

⁷ INAF-Istituto di Radioastronomia, Via P. Gobetti 101, I-40129 Bologna, Italy

⁸ Obserwatorium Astronomiczne Uniwersytetu Jagiellońskiego, ul. Orła 171, 30-244 Kraków, Poland

⁹ Department of Earth and Space Sciences, Chalmers University of Technology, Onsala Space Observatory, SE-43992 Onsala, Sweden

¹⁰ GEPI, Observatoire de Paris, UMR 8111, CNRS, Université Paris Diderot, 5 Place Jules Janssen, F-92190 Meudon, France

¹¹ Department of Physical Sciences, The Open University, Milton Keynes MK7 6AA, UK

¹² Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

¹³ Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands

¹⁴ Astrophysics, Department of Physics, Keble Road, Oxford OX1 3RH, UK

¹⁵ Department of Physics, University of the Western Cape, Private Bag X17, Bellville 7535, South Africa

¹⁶ School of Physics and Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK

¹⁷ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

¹⁸ RALSpace, The Rutherford Appleton Laboratory, Chilton, Didcot, Oxfordshire OX11 0NL, England

Received 2014 September 18; accepted 2014 October 6; published 2014 October 28

ABSTRACT

Carbon radio recombination lines (RRLs) at low frequencies ($\lesssim 500$ MHz) trace the cold, diffuse phase of the interstellar medium, which is otherwise difficult to observe. We present the detection of carbon RRLs in absorption in M82 with the Low Frequency Array in the frequency range of 48–64 MHz. This is the first extragalactic detection of RRLs from a species other than hydrogen, and below 1 GHz. Since the carbon RRLs are not detected individually, we cross-correlated the observed spectrum with a template spectrum of carbon RRLs to determine a radial velocity of 219 km s^{-1} . Using this radial velocity, we stack 22 carbon- α transitions from quantum levels $n = 468$ –508 to achieve an 8.5σ detection. The absorption line profile exhibits a narrow feature with peak optical depth of 3×10^{-3} and FWHM of 31 km s^{-1} . Closer inspection suggests that the narrow feature is superimposed on a broad, shallow component. The total line profile appears to be correlated with the 21 cm H I line profile reconstructed from H I absorption in the direction of supernova remnants in the nucleus. The narrow width and centroid velocity of the feature suggests that it is associated with the nuclear starburst region. It is therefore likely that the carbon RRLs are associated with cold atomic gas in the direction of the nucleus of M82.

Key words: galaxies: individual (M82) – galaxies: ISM – ISM: general – radio lines: galaxies – radio lines: ISM

Online-only material: color figures

1. INTRODUCTION

The nearby (3.52 ± 0.02 Mpc; Jacobs et al. 2009) nuclear starburst galaxy M82 has been observed to host a wide range of phases of the interstellar medium (ISM). Observations of disrupted H I within the disk show that the neutral gas is more concentrated in the nuclear region (Yun et al. 1993). There is a rotating ring of molecular gas in the nucleus, as seen from observations of HCN, HCO⁺, CO(2–1) (Kepley et al. 2014), OH (e.g., Argo et al. 2010), and CO images (e.g., Westmoquette et al. 2013). Numerous H II regions are seen (e.g., McDonald et al. 2002; Gandhi et al. 2011) in this area. The nuclear region is also studded with compact, bright supernova remnants (SNRs; e.g., Muxlow et al. 1994; Fenech et al. 2010). The spectral turnovers of the SNRs (Varenius et al. 2014; Wills et al. 1997) as well as the overall spectrum (Varenius et al. 2014; Adebahr et al. 2013) indicate the presence of free–free absorption by

ionized gas. The complex interplay of all these components of the ISM is not fully understood.

Carbon radio recombination lines (RRLs) can help characterize the cold, diffuse phase of the ISM. When free electrons recombine with atoms at quantum numbers $n \gtrsim 50$, the decreased energy spacing of subsequent levels produces RRLs. By comparing observations of RRLs with detailed physical models, we can determine information on the physical properties, such as electron temperature and density, of the gas in which the RRLs originate (e.g., F. Salgado et al., in preparation; Walmsley & Watson 1982; Shaver 1975; Dupree 1969). At frequencies $\lesssim 500$ MHz, RRLs are spaced closely enough that wide-bandwidth instruments like the Low Frequency Array (LOFAR; van Haarlem et al. 2013) are able to track the dependence of line properties on quantum number within a single observation. These properties make RRLs a powerful tool for determining the temperature and density of their host phases of the ISM.

RRLs fall into two categories: discrete and diffuse. Discrete RRLs trace warm ($T_e \sim 10^4$ K), high-density ($n_e > 100 \text{ cm}^{-3}$) gas associated with H II regions (Palmer 1967). They are predominantly seen at frequencies above ~ 1 GHz and originate from hydrogen, helium, and carbon (e.g., Poppi et al. 2007; Konovalenko & Stepkin 2005; Roelfsema & Goss 1991). These types of RRLs have been detected in a handful of nearby bright star-forming galaxies (e.g., Shaver et al. 1977; Anantharamaiah et al. 1993; Rodriguez-Rico et al. 2004; Roy et al. 2008).

Diffuse carbon RRLs (CRRLs) trace the cold neutral medium (CNM), which is cold ($T_e \sim 100$ K) and diffuse ($n_e \lesssim 0.1 \text{ cm}^{-3}$). Diffuse RRLs are observed at frequencies below 1 GHz. Typically, the CNM has ionization levels that are too low to produce hydrogen and helium lines, and only CRRLs are observed. The ionization energy for atomic carbon is only 11.3 eV, lower than that of hydrogen, 13.6 eV. Some photons that can escape H II regions can therefore ionize carbon, and thus C II is expected to be the dominant state of carbon in the ISM. While discrete RRLs are used extensively to study star-forming regions (e.g., Anderson et al. 2011; Roelfsema et al. 1992), not much is yet known about the CNM associated with diffuse RRLs, even in our Galaxy. Low-frequency observations have shown that the CRRL emitting and absorbing gas is prevalent on scales of degrees along the Galactic plane (Erickson et al. 1995; Kantharia & Anantharamaiah 2001). Pinhole studies in the direction of H II regions (e.g., Golyntkin & Konovalenko 1991), SNRs (Cassiopeia A; e.g., Asgekar et al. 2013; Payne et al. 1989; Konovalenko & Sodin 1981), or bright background extragalactic sources (Oonk et al. 2014) have detected CRRLs on smaller scales. The new LOFAR (van Haarlem et al. 2013) has observed CRRLs in our Galaxy. Over the next several years, LOFAR will perform a CRRL survey of the Galactic plane, on scales from degrees down to several arcseconds, producing maps that will provide a comprehensive picture of the Galactic CNM by quantifying the average gas temperatures and densities, as well as abundances.

In this Letter, we present the first detection of extragalactic CRRLs, observed in M82 at frequencies near 60 MHz with LOFAR. This is the first extragalactic detection of diffuse RRLs, and the first extragalactic detection of RRLs from a species other than hydrogen. This detection opens up the possibility of tracing the evolution of the CNM through all stages of galaxy formation.

Section 2 outlines observations, data reduction, and imaging. Section 3 describes the extraction and processing of the spectra, cross-correlation of the overall spectrum to determine a velocity, and subband stacking to achieve a detection. Results are presented in Section 4. Discussion and conclusions follow in Sections 5 and 6, respectively.

2. OBSERVATIONS AND DATA REDUCTION

M82 was observed with a total on-source time of 5.0 hr on 2013 February 21 with the LOFAR low-band antenna (LBA; 10–80 MHz) stations, as part of early science (Cycle 0) data obtained for the LOFAR Survey of nearby galaxies (LC0_043, PI: R. Beck). We used 13 remote and 24 core stations for this observation, giving baselines between 90 m and 85 km. Each LBA station contains 96 dual polarization dipole antennas, 48 of which can be used simultaneously, and has a total radius of 87 m. We used the LBA_OUTER configuration, where the outermost more sparsely spaced 48 antennas in each station record data. This configuration has a reduced field of view compared to other configurations, and mitigates the problem of mutual coupling between closely spaced antennas, providing slightly increased sensitivity.

Each dual-polarization dipole antenna provides four linear correlation products which were used to reconstruct Stokes I images. We obtained complete frequency coverage between 30 and 78 MHz with 1 s time resolution. This bandwidth was divided in 244 subbands, each 0.1953 MHz wide, and further subdivided into 128 channels, resulting in channel widths ranging from 6 to 16 km s^{-1} . Using the total available bandwidth of 488 subbands and the multibeam capability of LOFAR, we simultaneously observed the calibrator source 3C196 with identical frequency coverage.

The calibrator data were processed by the observatory pipeline, starting by flagging radio frequency interference (RFI) with the AOFlagger (Offringa 2010). Typically, a few percent of the data were flagged due to RFI, consistent with a study of the LOFAR RFI environment (Offringa et al. 2013). The calibrator data were then averaged to 16 channels and 6 s before calibration with the BlackBoard Selfcal software system (Pandey et al. 2009). The target data were flagged with the AOFlagger, averaged to 6 s and 32 channels and the amplitude calibration obtained from 3C196 was applied. The velocity sampling of the individual subbands in target data after averaging ranges from 29 to 37 km s^{-1} .

For imaging we selected only the 24 LOFAR core stations, which all observe with the same clock, making correction for time delays unnecessary. The proximity of the stations to each other also mitigates the chance of observing the target through severely different ionospheric conditions. Image cubes of a region with an area of $6 \times 6 \text{ deg}^2$ centered on M82 were made with AWimager (Tasse et al. 2013), imaging and cleaning each channel individually. We used Briggs weighting (Briggs 1995) with a robust value of 0.0 to create images with resolutions ranging between $259 \times 340 \text{ arcsec}^2$ and $322 \times 425 \text{ arcsec}^2$. At this resolution, M82 is unresolved. Low-resolution imaging parameters were deliberately chosen to speed up the computationally expensive imaging process. LOFAR does not support Doppler tracking so we have Doppler-corrected the data post imaging. In total, we imaged 69 subbands in the range 50–64 MHz, centered on the peak response of the LBA (van Haarlem et al. 2013).

3. SPECTRAL PROCESSING

3.1. Individual Subband Processing

We extracted the spatially integrated spectra from elliptical apertures, where the appropriate elliptical dimensions in each subband were determined by fitting a two-dimensional Gaussian to the point source in the center and using the curve-of-growth (see Figure 1) to select an aperture size that captured the total flux density of the point source but did not include nearby background sources. We calculated the standard deviation for each subband and implemented automatic removal of subbands with excessively large standard deviations ($\geq 10 \times \sigma_{\text{median}}$), verified by visual inspection. Five subbands were removed from further processing. We inspected each subband for individual detections of lines, which were not seen. In order to increase the signal-to-noise ratio, we stacked CRRL α -transitions ($\Delta n = 1$, for $468 < n < 508$) to form an average line profile.

3.2. Measuring the Velocity/Redshift

Maximizing the signal-to-noise ratio of a stacked line profile requires precisely stacking the centers of each individual line, which requires an accurate radial velocity (redshift). While the systemic velocity of M82 has been measured as $210 \pm 20 \text{ km s}^{-1}$ relative to the local standard of rest (LSR), observations show




Figure 1. Demonstration of the spectral extraction process. The top left panel shows the center of an image obtained by averaging all channels maps within one subband, with a white contour marking the extraction aperture (diameter ~ 15 arcmin). The top center panel shows the curve-of-growth for the same subband used to determine the size of the aperture. The top right panel shows the optical depth standard deviation as a function of subband (frequency), with excessively noisy subbands filtered out. The bottom panel displays the normalized raw extracted spectrum for the same subband as the top left and center panels. A linear fit is overplotted as a blue dotted line, and observed and rest-frame ($z = 0.00073$) frequencies of the α -transition CRRL in this subband are labeled.

(A color version of this figure is available in the online journal.)




Figure 2. Cross-correlation values vs. redshift. Larger absolute values indicate a higher correlation between the template and observed spectra.

(A color version of this figure is available in the online journal.)

that even in the nuclear region there is a spread of ~ 200 km s $^{-1}$ in velocity (e.g., Wills et al. 1998; Kepley et al. 2014). We constrained the velocity for the CRRL absorbing gas by cross-correlating a spectral template of CRRLs (using Gaussian line profiles at known CRRL frequencies) with the entire observed spectrum over a range of redshifts. The template line peaks were normalized to a value of 3×10^{-3} . For the less certain line width parameter, we ran a series of models with different line widths, starting at $v = 15$ km s $^{-1}$ (to avoid cross-correlating with noise peaks). The magnitude of the cross-correlation peaks at $v = 210$ km s $^{-1}$ relative to the LSR (optical velocity), and we iterated with a template that only contains subbands for which rest-frame CRRLs would appear at this redshift. This reduces the amount of noise which is cross-correlated, very slightly improving the definition of the peak of the cross-correlation.

Figure 2 shows a clear peak in the cross-correlation at $v = 219$ km s $^{-1}$ ($z = 0.00073 \pm 0.00003$), consistent with

the systemic velocity of M82, and we used this to stack our lines. We did not find evidence for CRRL features at either of the velocities of the two secondary peaks, $v = 168$ km s $^{-1}$ and $v = 255$ km s $^{-1}$.

3.3. Reconstructing the Line Profile

Requiring that rest-frame CRRL frequencies be at least six channels away from the edge of a subband to avoid problems with noisy edge channels, we find that 23 subbands have α -transitions. We clipped the first and last three edge channels and converted to optical velocity using the rest frequency of the CRRL within the subband. After blanking ± 50 km s $^{-1}$ around the expected velocity of the CRRL, we fitted the continuum in each subband individually with a low (first or second) order polynomial. We also tried blanking different ranges around the line, from ± 50 to ± 250 km s $^{-1}$, without seeing substantial differences in the final spectrum. For one subband, any blanking left continuum only on one side of the line, so we did not include this subband in the final stack. Each subband was continuum subtracted. The final spectrum was constructed from the individual points of 22 subbands; see Figure 3.

The final stacked spectrum has a velocity sampling of ~ 1.5 km s $^{-1}$ within approximately ± 150 km s $^{-1}$ of the measured velocity (219 km s $^{-1}$, LSR). The measurement error of each point is equal to the standard deviation of the continuum in the subband from which the point originates. The weighted standard deviation of the continuum in the final spectrum within approximately ± 150 km s $^{-1}$ of the expected CRRL line center is $\sigma_\tau = 0.005$, which is approximately $\sqrt{22}$ smaller than the average noise in a subband.

Galactic CRRLs show peak optical depths of 10^{-3} – 10^{-4} (e.g., Kantharia & Anantharamaiah 2001; Konovalenko & Sodin 1981). If the CRRLs in M82 have similar peak optical depths then these would be within the noise of this spectrum. We therefore used a low-pass Savitzky–Golay filter (SGF; Savitzky & Golay 1964) which is a special case of a least-squares smoothing function that convolves the data with a filter whose shape is dependent on the polynomial order used for fitting.




Figure 3. Stacked spectrum with the expected CRRL feature at the measured systemic velocity of $v = 219 \text{ km s}^{-1}$. From top to bottom: the complete reconstructed spectrum; the portion of the spectrum with dense velocity coverage; the normalized density of points in the spectrum. The red dashed line in the top two panels represents the continuum.

(A color version of this figure is available in the online journal.)

Other filtering methods (see Figure 4) produced similar line profiles. After testing SGF filter widths from 15 to 60 data points for both first and second order polynomials, we selected an SGF with a width of 31 data points, which provided the flattest continuum. Within approximately $\pm 200 \text{ km s}^{-1}$ of the feature in the final, smoothed spectrum, the standard deviation of the continuum is $\sigma_\tau = 3.3 \times 10^{-4}$, in good agreement with our expectations from the raw spectrum noise and filter window size. We assessed this filtering method with modeled spectra and found that the integrated line profile is preserved within the errors even after introducing Gaussian noise.

Radial velocities within the nuclear region range from ~ 120 to 300 km s^{-1} . The centroid of the CRRL feature is therefore consistent with an absorption feature associated with the center of the nuclear region. The cross-correlation is already strong evidence that the feature does not arise merely from a chance alignment of noise, since the feature only arises from a location where we know there to be gas, which is necessary but not sufficient for CRRLs. To rule out other possibilities, we

conducted a series of tests. We used a “jack-knife” procedure where we re-stacked the subbands, each time leaving out an individual subband. None of the stacks produced a significant change to the absorption feature, confirming that the stacked spectrum is not dominated by one subband. Additionally, we stacked the subbands after introducing random velocity shifts to each subband before stacking, and were unable to produce any credible features. Spectra extracted from the background sky also do not show an absorption feature when stacked. From these tests, we conclude that our feature is real and associated with the M82 starburst galaxy.

We tried to detect CRRL β -transitions ($\Delta n = 2$), without success. This is not unexpected, as the CRRL β -transitions are expected to have integrated optical depths that are only 15%–30% that of the α -transitions (e.g., Stepkin et al. 2007; Payne et al. 1989). We did not find evidence for local CRRLs at $z = 0$. A search for hydrogen RRLs, both locally and at the redshift of M82, was also unproductive. This is unsurprising as these are believed to be very weak at these frequencies (e.g., F. Salgado et al., in preparation; Shaver 1975).

4. RESULTS

We stacked the spectra to search for CRRLs in M82 using our measured velocity, $v = 219 \text{ km s}^{-1}$, and detect carbon α -transition RRLs with a combined signal-to-noise ratio of 8.5σ in the filtered spectrum. The central absorption feature can be fitted by a four parameter Gaussian profile with a depth of $2.8^{+0.12}_{-0.10} \times 10^{-3}$, a FWHM of $30.6^{+2.3}_{-1.0} \text{ km s}^{-1}$, a center of $211.3^{+0.7}_{-0.5} \text{ km s}^{-1}$, and an additive offset of $-2^{+0.012}_{-0.072} \times 10^{-4}$ (see Figure 4).

5. DISCUSSION

Other absorption features have been observed in M82 with similarly narrow widths. Weiß et al. (2010) observed $p\text{-H}_2\text{O}$ absorption (in the far infrared) with a FWHM of 60 km s^{-1} , although this is offset from the CRRL absorption by $\sim 50 \text{ km s}^{-1}$. Additionally, a narrow component results if we




Figure 4. Top: different methods of smoothing the noise. The various methods provide similar line profiles. Bottom: the final spectrum, fitted with a Gaussian profile. (A color version of this figure is available in the online journal.)




Figure 5. Comparison with other tracers. The CRRL profile is plotted in black. Molecular emission lines (Kepley et al. 2014) are shown by the dashed lines and are inverted for easier comparison, and H I absorption in the direction of SNRs in M82 (Wills et al. 1998) is plotted with solid colored lines. The gold solid line is the total contribution to the H I absorption profile in the direction of all SNRs in M82, while the dark blue line is the contribution from only those SNRs relatively unaffected by free-free absorption at low frequencies. The bottom panel shows the central portion of the spectra, with only the H I absorption profiles overlaid on the CRRL spectrum.

(A color version of this figure is available in the online journal.)

limit the reconstruction of the H I absorption profile in the direction of SNRs in the nucleus of M82 to only the handful of bright SNRs which are relatively unaffected by free-free absorption at low frequencies. We constructed the H I absorption line profiles from measurements in Wills et al. (1998). We interpolated the spectra (mJy beam^{-1}) in the direction of each of the 26 SNRs which show H I absorption onto a velocity grid with 1 km s^{-1} resolution using a cubic spline function. Addition of the individual spectra produces a combined spectrum with information in the direction of SNRs. For the low-frequency spectrum, we cross matched the SNRs in Table 1 of Wills et al. (1998) with those still seen by LOFAR at 154 MHz (Varenius et al. 2014). The eight SNRs from Wills et al. (1998) that have measured H I absorption and are $> 1 \text{ mJy}$ at 154 MHz are listed in that paper as $39.10+57.3$, $39.40+56.1$, $39.77+56.9$, $42.53+61.9$, $43.31+59.2$, $45.74+65.2$, $45.89+63.8$, and $46.52+63.9$.

Both of these absorption profiles show broader, shallower structure to either side of a deep, central feature, of which we see hints in the reconstructed CRRL profile (see Figure 5). Galactic observations show a correlation between CRRLs and H I (e.g., Kantharia & Anantharamaiah 2001; Payne et al. 1989), so it is unsurprising that this correlation would also be present in M82.

As Figure 5 illustrates, integration of other tracers of interstellar gas over the disk of M82 leads to FWHM in excess of 200 km s^{-1} (Kepley et al. 2014; Wills et al. 1998). This leads us to expect a broad component of the CRRL profile, although the signal to noise coupled with the fixed velocity coverage and location of CRRLs in the subbands makes this difficult to ascertain. The overall CRRL profile follows the low-frequency H I line profile (Figure 5). This shows that the CRRL absorbing gas is preferentially observed in front of those continuum sources that are still bright at low frequencies. The continuum offset and low-level absorption features in the CRRL spectrum also show reasonable agreement with the H I absorbing gas, indicating that possibly we are observing a widespread CRRL component associated with cold, diffuse H I throughout M82.

Integrating the line profile gives $\int \tau_{\text{CRRL}} dv = 21.3 \text{ s}^{-1}$. To constrain physical parameters of the gas, i.e., temperature and

density, a single measurement is not enough. Additional measurements at higher frequencies (100–400 MHz) are necessary.

6. CONCLUSIONS

We have presented the first detection of extragalactic CRRLs, in M82, using early science (Cycle 0) observations with LOFAR. While we already know that cold, neutral gas exists in M82 from H I measurements, this is the first independent measurement of the atomic CNM. The narrow CRRL line is at a velocity corresponding to the center of M82 and the profile corresponds to the absorption feature seen in the H I spectrum. Higher resolution, higher sensitivity studies of M82 at the same frequencies would help to confirm that this feature is indeed associated with the nuclear region, and potentially shed more light on the origin of the optical depth. To constrain the gas properties we need studies at higher frequencies, and we are working toward detecting CRRLs in the range of 120–240 MHz with LOFAR’s high band antenna array.

This discovery paves the way for future extragalactic CRRL studies to trace the CNM throughout the formation and evolution of galaxies, and is the basis of a pilot survey for CRRLs in other extragalactic sources.

L.K.M. acknowledges financial support from NWO Top LOFAR project, project No. 614.001.006. J.B.R.O. acknowledges financial support from NWO Top LOFAR-CRRL project, project No. 614.001.351. The authors thank A. Kepley for providing GBT observed line profiles.

REFERENCES

- Adebahr, B., Krause, M., Klein, U., et al. 2013, *A&A*, **555**, A23
- Anantharamaiah, K. R., Zhao, J.-H., Goss, W. M., & Viallefond, F. 1993, *ApJ*, **419**, 585
- Anderson, L. D., Bania, T. M., Balsler, D. S., & Rood, R. T. 2011, *ApJS*, **194**, 32
- Argo, M. K., Pedlar, A., Beswick, R. J., Muxlow, T. W. B., & Fenech, D. M. 2010, *MNRAS*, **402**, 2703
- Asgekar, A., Oonk, J. B. R., Yatawatta, S., et al. 2013, *A&A*, **551**, L11

- Briggs, D. S. 1995, *BAAS*, 27, 112.02
- Dupree, A. K. 1969, *ApJ*, 158, 491
- Erickson, W. C., McConnell, D., & Anantharamaiah, K. R. 1995, *ApJ*, 454, 125
- Fenech, D., Beswick, R., Muxlow, T. W. B., Pedlar, A., & Argo, M. K. 2010, *MNRAS*, 408, 607
- Gandhi, P., Isobe, N., Birkinshaw, M., et al. 2011, *PASJ*, 63, 505
- Golynkin, A. A., & Konovalenko, A. A. 1991, *SvAL*, 17, 7
- Jacobs, B. A., Rizzi, L., Tully, R. B., et al. 2009, *AJ*, 138, 332
- Kantharia, N. G., & Anantharamaiah, K. R. 2001, *JApA*, 22, 51
- Kepley, A. A., Leroy, A. K., Frayer, D., et al. 2014, *ApJL*, 780, L13
- Konovalenko, A. A., & Sodin, L. G. 1981, *Natur*, 294, 135
- Konovalenko, A. A., & Stepkin, S. V. 2005, in *Radio Recombination Lines*, ed. L. I. Gurvits, S. Frey, & S. Rawlings (EAS SP-15; Cambridge: Cambridge Univ. Press), 271
- McDonald, A. R., Muxlow, T. W. B., Wills, K. A., Pedlar, A., & Beswick, R. J. 2002, *MNRAS*, 334, 912
- Muxlow, T. W. B., Pedlar, A., Wilkinson, P. N., et al. 1994, *MNRAS*, 266, 455
- Offringa, A. R. 2010, *ascl soft*, 10017
- Offringa, A. R., de Bruyn, A. G., Zaroubi, S., et al. 2013, *A&A*, 549, A11
- Oonk, J. B. R., van Weeren, R. J., Salgado, F., et al. 2014, *MNRAS*, 437, 3506
- Palmer, P. 1967, *ApJ*, 149, 715
- Pandey, V. N., van Zwieten, J. E., de Bruyn, A. G., & Nijboer, R. 2009, in *ASP Conf. Ser. 407, The Low-Frequency Radio Universe*, ed. D. J. Saikia, D. A. Green, Y. Gupta, & T. Venturi (San Francisco, CA: ASP), 384
- Payne, H. E., Anantharamaiah, K. R., & Erickson, W. C. 1989, *ApJ*, 341, 890
- Poppi, S., Tsivilev, A. P., Cortiglioni, S., Palumbo, G. G. C., & Sorochenko, R. L. 2007, *A&A*, 464, 995
- Rodriguez-Rico, C. A., Viallefond, F., Zhao, J.-H., Goss, W. M., & Anantharamaiah, K. R. 2004, *ApJ*, 616, 783
- Roelfsema, P. R., & Goss, W. M. 1991, *A&AS*, 87, 177
- Roelfsema, P. R., Goss, W. M., & Mallik, D. C. V. 1992, *ApJ*, 394, 188
- Roy, A. L., Goss, W. M., & Anantharamaiah, K. R. 2008, *A&A*, 483, 79
- Savitzky, A., & Golay, M. J. E. 1964, *AnaCh*, 36, 1627
- Shaver, P. A. 1975, *A&A*, 43, 465
- Shaver, P. A., Churchwell, E., & Rots, A. H. 1977, *A&A*, 55, 435
- Stepkin, S. V., Konovalenko, A. A., Kantharia, N. G., & Udaya Shankar, N. 2007, *MNRAS*, 374, 852
- Tasse, C., van der Tol, S., van Zwieten, J., van Diepen, G., & Bhatnagar, S. 2013, *A&A*, 553, A105
- van Haarlem, M. P., Wise, M. W., Gunst, A. W., et al. 2013, *A&A*, 556, A2
- Varenius, E., Conway, J. E., Martí-Vidal, I., et al. 2014, *A&A*, submitted
- Walmsley, C. M., & Watson, W. D. 1982, *ApJ*, 260, 317
- Weiß, A., Requena-Torres, M. A., Güsten, R., et al. 2010, *A&A*, 521, L1
- Westmoquette, M. S., Smith, L. J., Gallagher, J. S., & Walter, F. 2013, *MNRAS*, 428, 1743
- Wills, K. A., Pedlar, A., & Muxlow, T. W. B. 1998, *MNRAS*, 298, 347
- Wills, K. A., Pedlar, A., Muxlow, T. W. B., & Wilkinson, P. N. 1997, *MNRAS*, 291, 517
- Yun, M. S., Ho, P. T. P., & Lo, K. Y. 1993, *ApJL*, 411, L17