

Depletion of Intense Fields

D. Seipt,^{1,*} T. Heinzl,² M. Marklund,³ and S. S. Bulanov⁴

¹*Helmholtz-Institut Jena, Fröbelstieg 3, 07743 Jena, Germany*

²*School of Computing, Electronics and Mathematics, Plymouth University, Plymouth PL4 8AA, United Kingdom*

³*Department of Physics, Chalmers University of Technology, SE-41296 Gothenburg, Sweden*

⁴*Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

(Received 2 May 2016; published 14 April 2017)

The interaction of charged particles and photons with intense electromagnetic fields gives rise to multiphoton Compton and Breit-Wheeler processes. These are usually described in the framework of the external field approximation, where the electromagnetic field is assumed to have infinite energy. However, the multiphoton nature of these processes implies the absorption of a significant number of photons, which scales as the external field amplitude cubed. As a result, the interaction of a highly charged electron bunch with an intense laser pulse can lead to significant depletion of the laser pulse energy, thus rendering the external field approximation invalid. We provide relevant estimates for this depletion and find it to become important in the interaction between fields of amplitude $a_0 \sim 10^3$ and electron bunches with charges of the order of 10 nC.

DOI: 10.1103/PhysRevLett.118.154803

The interaction of charged particles with ultraintense electromagnetic (EM) pulses is the cornerstone of a newly emerging area of research, high intensity particle physics, located at the intersection of quantum electrodynamics (QED) and the theory of strong EM background fields. The latter significantly alter the physics of typical QED processes, leading to effects not encountered in perturbative quantum field theory [1–6]. Recently, there has been a surge of interest in these processes due to the planning and realization of new laser facilities, which will be able to deliver EM pulses of unprecedented intensities to test the predictions of high intensity particle physics [2]. Moreover, the development of compact multi-GeV laser electron accelerators [1,2,7,8] adds another component necessary to carry out these studies.

Here, we will assume that the strong EM field is provided by an ultraintense laser (pulse) with wave vector k , central frequency $\omega = 2\pi/\lambda$ in the optical regime, and electric field magnitude E . The interactions of this strong field with photons and charged particles are parametrized in terms of the following parameters (we use natural units throughout, $\hbar = c = 1$): (i) the (Lorentz and gauge invariant [9]) dimensionless amplitude of the EM vector potential, $a_0 = eE/\omega m$; (ii) the QED critical field, $E_S = m^2/e$ [10]; (iii) the strong field invariants $\chi_e^2 = -e^2(F^{\mu\nu}p_\nu)^2/m^6$ and $\chi_\gamma^2 = -e^2(F^{\mu\nu}k'_\nu)^2/m^6$ [6]. Here, e and m are electron charge and mass, $F_{\mu\nu}$ is the EM field tensor, while p_ν and k'_ν denote the four momenta of electron and photon probing the laser. The parameter a_0 is usually referred to as the classical nonlinearity parameter, since its physical meaning is the energy gain of an electron (in units m) traversing a reduced wavelength $\lambda = 1/\omega$ of the field. For $a_0 > 1$ the electron or positron motion in such a field becomes relativistic.

The parameter E_S characterizes a distinct feature of QED, the ability to produce new particles from vacuum. This happens when an energy of mc^2 is delivered across an electron Compton wavelength $\lambda_e = 1/m$, which is precisely achieved by E_S [10]. The parameters χ_e and χ_γ characterize the interaction of charged particles and photons with the strong EM field. For example, χ_e is the EM field strength in the electron rest frame in units of E_S . Quantum effects become of crucial importance when $E \approx E_S$ or $\chi_{e,\gamma} \sim 1$.

For large field amplitudes $a_0 \gg 1$, the interaction of electrons or positrons and photons with strong EM fields involves the absorption of a large number of photons from the field. Clearly, these correspond to an energy loss of the laser background field, which may or may not be negligible. Revisiting the results on multiphoton Compton and Breit-Wheeler processes [5,6,11], we find that there is indeed a parameter range, for which the depletion of the laser becomes substantial. The processes in question have recently received a lot of interest [12,13], albeit with a focus on the *final* states (a frequency shifted photon or electron positron pairs).

In this Letter, we want to change perspective and study in detail the dependence of nonlinear Compton scattering on the *initial* multiphoton states, that is on the number of laser photons absorbed. This will allow us to establish a threshold for the validity of the external field approximation and discuss some immediate consequences. These findings should have a direct impact on the analysis of QED backreaction on the classical EM field [14]. It should also be of great importance for the study of EM avalanches [15–17], since background depletion will significantly alter the energy partitioning of the processes. An avalanche is formed when Compton and Breit-Wheeler processes occur subsequently in an EM field of sufficiently high intensity,

resulting in an exponential growth of the number of emitted particles.

The external field approximation is valid when the number of photons absorbed from the laser ΔN_A is small compared to the total number $N_L \gg 1$ of photons in the pulse, which we take to be focused to volume $V = \lambda^3$. A natural criterion for depletion is then provided by the equality $\Delta N_A = N_L$. The number of laser photons is proportional to intensity or field strength squared, $N_L \approx (2\pi/\alpha)(\lambda^2/\lambda_e^2)a_0^2 \approx 2 \times 10^{14} a_0^2$. Here, $\alpha = e^2/4\pi \approx 1/137$ denotes the fine structure constant. The number of absorbed photons is $\Delta N_A \approx (\Delta E/\omega)N_T$, where N_T is the number of electrons in the bunch and $\Delta E = P_{\text{rad}}T$ is the energy loss upon radiating power P_{rad} per laser period T . This power, a Lorentz invariant, can be estimated classically by making an analogy with synchrotron radiation [18,19]. In consequence, we will be able to estimate the number of photons absorbed from the field, the characteristic energy of an emitted photon, and the angle of emission, implying a rather complete characterization of the processes. To this end we go to a boosted frame, where the electron is on average at rest. If the laser is circularly polarized, the electron moves on a circle like in a synchrotron with four velocity $u = \gamma(1, \boldsymbol{\beta}_\perp, 0)$, where $\gamma^2 = (1 - \boldsymbol{\beta}_\perp^2)^{-1} = 1 + a_0^2$ characterizes the average rest frame (ARF). Using Larmor's formula, the radiated power becomes $P_{\text{rad}} = -(2/3)\alpha i u^2 = (2/3)\alpha \omega^2 a_0^2(1 + a_0^2)$. The boost to the ARF may be realized by choosing the initial electron momentum $p = m\gamma_e(1, 0, 0, -\beta_e)$, such that its light-front component equals $p^- = m\gamma_e(1 + \beta_e) = m(1 + a_0^2)^{1/2} \equiv m_*$, with m_* denoting the intensity-dependent effective mass [20]. [If ℓ is an arbitrary four vector its scalar product with the laser momentum can be written as $k \cdot \ell = \omega(\ell^0 - \ell_z) \equiv \omega \ell^-$, which defines the light-front component ℓ^- [21].] At high energy $\gamma_e \gg 1$, the radiation is emitted in the plane of electron motion, which in the ARF is perpendicular to the laser axis. In the lab frame this transforms into an emission angle

$$\tan \theta = P_\perp/P_z = (2a_0 m/p^-)/[(m_*/p^-)^2 - 1] \quad (1)$$

determined by the ratio of longitudinal and transverse momenta in the ARF. For $a_0 \ll 1$ the emission angle is $\sim 1/\gamma_e$, hence small, while for $a_0 \gg 1$ there is significant emission in the transverse direction. In the ARF, $\theta = \pi/2$. The number of absorbed photons per laser period $T = 2\pi/\omega$ is then

$$\Delta N_A = (4\pi/3)\alpha a_0^2(1 + a_0^2)N_T. \quad (2)$$

So for $a_0 \gg 1$, the radiated power, hence the number of absorbed photons per laser cycle, increases like a_0^4 . From synchrotron radiation it is known that the power radiated into the s th harmonic asymptotically scales like $P_s \sim s^{1/3}$ [19], so that the total power is $P = \sum_s^{s_0} P_s \sim s_0^{4/3} \sim a_0^4$. We thus obtain the important result that the typical number s_0 of laser

photons, absorbed to yield emission of a single high-energy photon, scales like $s_0 \sim a_0^3$.

Turning back to the question of beam depletion, we equate $\Delta N_A \sim N_L$ to see that depletion requires

$$a_0^2 N_T \sim 6.5 \times 10^{15}. \quad (3)$$

For an electron bunch containing a charge of 1 nC, a laser with $a_0 \approx 10^3$ is needed. For such values of a_0 the energy ω' of the emitted photons is of the order of the electron energy gain per laser period, and the emission angle significantly deviates from $\sim 1/\gamma_e$. Thus, in this case, one expects not just a significant radiation reaction with ensuing changes of the particle trajectories [22] but also a strong recoil of the electron momentum. These features are best described in quantum theory, to which we now turn.

First, we estimate the depletion threshold by taking into account the discrete nature of photon emission. While the average number of absorbed photons s_0 still follows the classical scaling law $s_0 \sim a_0^3$ for $a_0 \gg 1$ [6], the classical formula (2) is replaced by

$$\Delta N_A \sim s_0 N_T (\lambda/L_C) \sim a_0^3 N_T (\lambda/L_C). \quad (4)$$

Here, L_C is the radiation length of the electron in a strong EM field [6] so that, on average, there is one photon emission per distance L_C . The classical behavior (2) is recovered in the limit $\chi_e \ll 1$ where $L_C \sim \lambda/a_0$. In the deep quantum regime $\chi_e \gg 1$, we employ the asymptotic expression $L_C = 0.43\lambda\gamma_e^{1/3}a_0^{-2/3}$ [6] to obtain a quantum formula for the threshold of depletion,

$$a_0^2 N_T / (a_0 \gamma_e)^{1/3} \sim 10^{14}, \quad (5)$$

which supersedes (3). Again, for an electron beam of 1 nC and $\gamma_e \approx 10^3$, a laser with intensity $a_0 \approx 10^3$ is required. Intensities of this magnitude should become a reality in the near future [2]. The critical value of a_0 depends weakly on the initial electron energy ($\sim \gamma_e^{1/5}$) as shown in Fig. 1, where $a_0(\gamma_e)$ is shown for different values of N_T . Thus, taking quantum effects into account increases the critical value of a_0 needed to deplete the laser pulse for a given value of initial electron momentum. We note that, for $\chi_e \gg 1$, the depletion of the electron beam energy is quite strong [23]. This corresponds to the threshold for depletion of the laser beam going down from (5) to (3), as illustrated in Fig. 1. Both (3) and (5) show that, when a sufficiently charged electron bunch collides with an intense laser pulse, depletion of the laser pulse can become significant, with the originally strong EM field turning weak. The required number of electrons is quite typical for an EM avalanche [16], where an intense laser produces a copious amount of high energy photons and subsequently electron-positron pairs. As the required electron densities are quite substantial, we briefly address the issue of coherence effects. To this end we note that the interelectronic distance




FIG. 1. The number N_T of radiating electrons required to see significant depletion plotted as a function of γ_e and a_0 . Black curves indicate the depletion thresholds $a_0(\gamma_e)$ when $N_T = 10^9$, 10^{10} , and 10^{11} . The dashed white curves represent $\omega'\Delta x = \text{const} \gg 1$, so that photon emission is incoherent across the relevant parameter space. Left, using (4). Right, using (6) below.

$\Delta x \sim \lambda/\sqrt{3}N_T$ remains much larger than the emitted photon wavelength $\propto 1/\omega'$. As a result, $\omega'\Delta x \gg 1$ (see Fig. 1), a well-established criterion for hard photon emission to be *incoherent* [24].

Second, we refine the depletion threshold estimate by calculating the quantum corrections to the average number of absorbed photons, taking into account the probabilistic nature of photon emission. In line with the current understanding of high-intensity laser matter interactions in the quantum regime, we model photon emission as a succession of incoherent one-photon events [17,25–27]. There will be regimes where this assumption becomes challenged, for instance when extreme field strengths are reached such that $\alpha\chi^{2/3} \gtrsim 1$. In this case, higher-order diagrams such as self-energy corrections [28] and coherent multiphoton emission [29] can no longer be neglected. A detailed investigation of higher-order effects is clearly beyond the scope of the present study, but we can at least state that $\alpha\chi^{2/3} \lesssim 0.1$ for our parameter range. It is thus sufficient to introduce one-particle emission probabilities $dP^{e,\gamma}/ds$, which are differential in the number of photons s absorbed from the laser field. The average amount of energy $\langle \mathcal{E} \rangle$ drawn from the laser field in a single photon emission or pair production is then $\langle \mathcal{E} \rangle = \omega\langle s \rangle$, with the average number of absorbed laser photons given by the expectation value $\langle s \rangle_{e,\gamma} = Z^{-1} \int ds s (dP^{e,\gamma}/ds)$ with normalization integral $Z = \int ds dP^{e,\gamma}/ds$.

In a monochromatic plane wave laser field, taken to be circularly polarized for simplicity, the variable s is discrete and describes the emission of higher harmonics due to absorption of s laser photons. Introducing the usual *quasimomentum* $q = p + (m^2 a_0^2 / 2k \cdot p)k$ (and analogously for q' , whence $q^2 = q'^2 = m_*^2$), kinematics become encoded in quasimomentum conservation, $q + sk = q' + k'$. The partial probabilities (per unit time) P_s^e were

calculated long ago [6,11] and give the total probability for Compton photon emission when summed over all harmonics: $P^e = \sum_{s=1}^{\infty} P_s^e$. For large values of $a_0 \gg 1$, the number s of harmonics contributing grows like $s \sim a_0^3$, and hence it can be assumed quasicontinuous. The sum may thus be replaced by an integral over s with integrand [6]

$$\frac{dP^e}{ds} = \frac{4\alpha\omega s}{1+a_0^2} \left(\frac{2}{s}\right)^{2/3} \int_0^1 \frac{dt}{(1+su_1t)^2} \left\{ -\Phi^2(\eta) + a_0^2 \left(\frac{2}{s}\right)^{2/3} \times \left(1 + \frac{s^2 u_1^2 t^2}{2(1+su_1t)}\right) [\eta\Phi^2(\eta) + \Phi'(\eta)^2] \right\}. \quad (6)$$

Φ and Φ' denote the Airy function and its derivative, their argument being $\eta(s,t) = (s/2)^{2/3} [1 - 4a_0^2(m/m_*)^2 t(1-t)]$, with $t = u/(su_1)$, $u = (k' \cdot k)/(p' \cdot k)$, and $u_1 = 2(k \cdot p)/m_*^2$. The dependence of $s dP^e/ds$ on s is shown in Fig. 2(b). The maximum corresponding to the most probable number of absorbed photons shifts towards lower values of s with an increase of initial electron energy. In Fig. 2(a) the dependence of $\langle s \rangle_e$ on the parameter a_0 shows an increase of the number of absorbed photons with the EM field strength, but indicates a dependence different from the classical behavior $s_0 \sim a_0^3$: the most probable number of absorbed photons also depends on the parameter χ_e as given by the fit $\langle s \rangle_e = 0.54a_0^3/(1 + 1.49\chi_e^{0.59})$. Using $\chi_e = 2\gamma_e a_0 \omega/m$, the threshold for depletion becomes $a_0^{1.08} \gamma_e^{-0.92} N_T \sim 6.8 \times 10^{11}$. For instance, when $a_0 = 10^3$, we find a value of $N_T \approx 10^{11-12}$, cf. Fig. 1 (right), which is larger than $N_T \approx 10^{10}$ predicted by the simple estimate (5), but still within reach of EM avalanches [16].

We note that for classical synchrotron emission it is straightforward to relate radiated to absorbed power, because of the continuity of emission. In the quantum case a typical




FIG. 2. The dependence of the average number $\langle s \rangle_e$ of absorbed laser photons on the dimensionless amplitude a_0 of the EM field for different values of γ_e . The corresponding distributions $s dP^e/ds$ for $a_0 = 200$ are shown in the inset (b). Black dotted curves represent the numerical fit provided.

interaction of an electron beam with an intense laser pulse proceeds via multiple emissions, each of them potentially resulting in a significant change of the electron momentum. To characterize such interactions one uses simulation codes with QED modules, which take into account multiphoton Compton and Breit-Wheeler processes. For these codes to be used for depletion calculations, each Compton process needs to be characterized by photon energy, angle of emission, and the number of absorbed photons. Furthermore, in numerical (QED-particle-in-cell) simulations of multistage emission processes, which lead to the formation of avalanches or cascades, photon and electron emission angles strongly determine the probability of the subsequent pair production or photon emission process, respectively [4,17,25,27,30,31]. We hence proceed by calculating these quantities.

The probabilities dP^e/ds determine the number distribution of photons absorbed from the laser field in a single high-frequency photon emission. In what follows, we relate dP^e/ds to the distribution $dP^e/d\chi_\gamma$ of the scattered photon longitudinal momentum ($\chi_\gamma \sim k \cdot k'$), which determines the intensity of the emitted high-energy photon radiation, via the chain rule:

$$\frac{dP^e}{d\chi_\gamma} = \frac{ds(\chi_\gamma)}{d\chi_\gamma} \frac{dP^e}{ds}, \quad (7)$$

where the functional relation $s = s(\chi_\gamma)$ is unknown. From the t integral in (6) we see that the integrand is sharply peaked at $t = 1/2$. Using energy momentum conservation we can solve $t = u/(su_1) = 1/2$ for s with the result

$$s(\chi_\gamma) = \frac{a_0^3}{\chi_e \chi_e - \chi_\gamma}. \quad (8)$$

This is valid for $a_0 \gg 1$ and reproduces the leading order of the related result (18) in [32]. A direct *numerical* determination of $s(\chi_\gamma)$ from (7) shows an excellent agreement with (8) for the most important range of $s \sim a_0^3$ (but deviates for $s \rightarrow 0$). Thus, when a Compton photon with a given value of χ_γ is emitted, the number of laser photons drawn from the laser field can safely be estimated using (8) within the model of one-photon incoherent emission. A formula completely analogous to (8) holds for the Breit-Wheeler process, $\gamma + s\gamma_L \rightarrow e^+e^-$, which becomes possible above a threshold in photon number, $s \geq s_0 = 2a_0(1 + a_0^2)/\chi_\gamma$. Details will be discussed elsewhere.

To further illustrate the power of the result (8), we employ it to determine the most probable emission angle without referring to an angular probability distribution. Let us write the scattered photon momentum as $k' = (\omega', \mathbf{k}'_\perp, k'_z)$, where $k'^2 = 0$. We can then find k'_\perp from quasimomentum conservation. Assuming a head-on collision of electrons and laser ($\mathbf{p}_\perp = 0$) the following answer is obtained:

$$k'^2_\perp = 2sk \cdot k' - \left(\frac{k \cdot k'}{k \cdot p}\right)^2 (m_*^2 + 2sk \cdot p). \quad (9)$$

This identity is manifestly invariant with respect to boosts collinear with the laser direction k . It defines an ellipse in the




FIG. 3. The distribution $s dP^e/ds$ as a function of k'_\perp and k'_z for $\gamma_e = 2000$, and for $a_0 = 200$ (left) and 2000 (right). The distributions are supported on a line $k'_\perp/k'_z = 1/10$ and $4/3$ in the left and right panels, respectively, cf. (10). Full curves on the floor represent the ellipses (9) for different values of s .

(k'_z, k'_\perp) plane for given values of γ_e , s , and a_0 , see Fig. 3. Plugging (8) into (9) yields the tangent of the most probable photon emission angle,

$$\tan \theta = k'_\perp/k'_z = \frac{4a_0\gamma_e}{4\gamma_e^2 - a_0^2}, \quad (10)$$

where $a_0, \gamma_e \gg 1$. This coincides with the classical emission angle (1) and is indeed consistent with the findings of [13]: As long as $\gamma_e \gg a_0$, the photons are predominantly emitted in the forward direction, with $\theta \sim a_0/\gamma_e \ll 1$. However, as a_0 increases, significant photon emission takes place in the perpendicular direction. This can be understood classically, in particular in the ARF, where $a_0 \approx 2\gamma_e$ ($p^- = m_*$), so that $\theta = \pi/2$ as required for circular (synchrotron) motion in the transverse plane as well as by (10). Equivalently, this follows from the classical equation of motion by calculating $\tan \theta = (\pi_\perp/\pi_z)_{\text{rms}}$, the ratio of the rms values of the classical electron momentum components in the laser field A , $\pi^\mu = p^\mu - eA^\mu + (ep \cdot A - e^2 A^2/2)k^\mu/k \cdot p$.

Going back to Fig. 3 we see that the distribution of emitted photons is essentially supported on a straight line, $k'_\perp/k'_z = \text{const}$ (with an angular spread of the order $1/a_0 \ll 1$), which intersects the ellipse (9) in a single point. To relate back to the topic of depletion we recall Fig. 1 (right), which tells us that we have to stay away from the axes and the origin in the $a_0 - \gamma_e$ plane according to our assumption of incoherent emission. The “safe” regime is thus $a_0 \sim \gamma_e \gg 1$, so that in terms of the emission angle we need to stay away from collinear emission, $a_0 \ll \gamma_e$ or $a_0 \gg \gamma_e$. Thus, in the generic regime of interest $a_0 \sim \gamma_e \gg 1$, there is substantial transverse emission, cf. Fig. 3, right, for which the emission angle is about 50° , with a depletion threshold of $N_T \approx 10^{10}$ according to Fig. 1.

In this Letter, we have reconsidered the multiphoton Compton process in strong EM fields, focusing on the energy loss of the laser due to absorption, which transforms the initially strong fields into weak ones. We found that this phenomenon has an intensity threshold of $a_0 \sim 10^3$, and requires $N_T \gtrsim 6.8 \times 10^{11} \gamma_e^{0.92} a_0^{-1.08}$ electrons per laser wavelength cubed, according to the numerical fit in Fig. 2. We have neglected coherent photon emission, which is

valid when $a_0 \sim \gamma_e \gg 1$. It is expected that the depletion threshold will be overcome in the case of EM avalanches. Thus, laser depletion will not just be due to pair creation as considered previously, but must also be taken into account in laser photon absorption.

We have further analyzed the photon emission rates differential in multiphoton number s and discovered that they strongly peak at a value s_0 , recall (8), which determines the direction of the photon emission relative to the initial electron momentum direction in terms of an emission angle θ via (10). For generic depletion parameters $a_0 \sim \gamma_e \gg 1$, one finds substantial emission in the transverse direction. In the collinear regime $\theta \ll 1$ (forward scattering, $a_0 \ll \gamma_e$) and $\theta \approx \pi$ (backscattering, $a_0 \gg \gamma_e$), coherent emission can no longer be neglected. Backscattering should dominate in the EM avalanche regime, i.e., in colliding laser pulses or during interactions of laser pulses with solid density foils or plasmas of near-critical density. The classical interpretation of the emission angle θ in terms of averages over trajectories should yield a new test of the particle-in-cell codes currently in use.

In future work, we want to understand the effect of depletion on the emission probabilities. This will require estimating the effect of a decreasing a_0 on, e.g., (6), building on previous work such as [33].

We acknowledge support from the Office of Science of the U.S. DOE under Contract No. DE-AC02-05CH11231. M.M. was supported by the Swedish Research Council Grants No. 2012-5644 and No. 2013-4248. We would like to thank Anton Ilderton for fruitful discussions. The authors acknowledge the hospitality of the Kavli Institute for Theoretical Physics (KITP), where this research was initiated during the Frontiers of Intense Laser Physics Program, and so was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915.

*Present address: Lancaster University, Physics Department, Bailrigg, Lancaster LA1 4YW, United Kingdom and Cockcroft Institute, Daresbury Laboratory, Keckwick Ln, Warrington WA4 4AD, United Kingdom.

†d.seipt@lancaster.ac.uk

- [1] G. Mourou, T. Tajima, and S. V. Bulanov, *Rev. Mod. Phys.* **78**, 309 (2006); M. Marklund and P. Shukla, *Rev. Mod. Phys.* **78**, 591 (2006); Y.I. Salamin, S. X. Hu, K. Z. Hatsagortsyan, and C. H. Keitel, *Phys. Rep.* **427**, 41 (2006); F. Ehlötzky, K. Krajewska, and J. Z. Kamiński, *Rep. Prog. Phys.* **72**, 046401 (2009); A. Di Piazza, C. Müller, K. Z. Hatsagortsyan, and C. H. Keitel, *Rev. Mod. Phys.* **84**, 1177 (2012).
- [2] M. Dunne, *Nat. Phys.* **2**, 2 (2006); *ELI-Extreme Light Infrastructure Science and Technology with Ultra-Intense Lasers WHITEBOOK*, edited by G. A. Mourou, G. Korn, W. Sandner, and J. L. Collier (THOSS Media GmbH, Berlin, 2011).
- [3] S. V. Bulanov *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **660**, 31 (2011); *AIP Conf. Proc.* **1507**, 825 (2012).
- [4] S. S. Bulanov, C. B. Schroeder, E. Esarey, and W. P. Leemans, *Phys. Rev. A* **87**, 062110 (2013).
- [5] H. Reiss, *J. Math. Phys. (N.Y.)* **3**, 59 (1962).
- [6] A. I. Nikishov and V. I. Ritus, *Sov. Phys. JETP* **19**, 529 (1964); **19**, 1191 (1964); V. I. Ritus, *J. Sov. Laser Res.* **6**, 497 (1985).
- [7] E. Esarey, C. B. Schroeder, and W. P. Leemans, *Rev. Mod. Phys.* **81**, 1229 (2009).
- [8] W. P. Leemans *et al.*, *Phys. Rev. Lett.* **113**, 245002 (2014).
- [9] T. Heinzl and A. Ilderton, *Opt. Commun.* **282**, 1879 (2009);
- [10] F. Sauter, *Z. Phys.* **69**, 742 (1931); W. Heisenberg and H. Euler, *Z. Phys.* **98**, 714 (1936); J. Schwinger, *Phys. Rev.* **82**, 664 (1951).
- [11] G. Breit and J. A. Wheeler, *Phys. Rev.* **46**, 1087 (1934); L. S. Brown and T. W. B. Kibble, *Phys. Rev.* **133**, A705 (1964); I. I. Goldman, *Sov. Phys. JETP* **19**, 954 (1964); V. P. Yakovlev, *Sov. Phys. JETP* **22**, 223 (1966).
- [12] N. B. Narozhnyi and M. S. Fofanov, *J. Exp. Theor. Phys.* **83**, 14 (1996); M. Boca and V. Florescu, *Phys. Rev. A* **80**, 053403 (2009); T. Heinzl, A. Ilderton, and M. Marklund, *Phys. Lett. B* **692**, 250 (2010); T. Heinzl, D. Seipt, and B. Kämpfer, *Phys. Rev. A* **81**, 022125 (2010); F. Mackenroth and A. Di Piazza, *Phys. Rev. A* **83**, 032106 (2011); A. I. Titov, H. Takabe, B. Kämpfer, and A. Hosaka, *Phys. Rev. Lett.* **108**, 240406 (2012); K. Krajewska and J. Z. Kamiński, *Phys. Rev. A* **85**, 062102 (2012).
- [13] C. Harvey, T. Heinzl, and A. Ilderton, *Phys. Rev. A* **79**, 063407 (2009); see also Ehlötzky *et al.*, Ref. [1].
- [14] F. Cooper and E. Mottola, *Phys. Rev. D* **40**, 456 (1989); Y. Kluger, J. M. Eisenberg, B. Svetitsky, F. Cooper, and E. Mottola, *Phys. Rev. Lett.* **67**, 2427 (1991); *Phys. Rev. D* **45**, 4659 (1992); N. B. Narozhny, S. S. Bulanov, V. D. Mur, and V. S. Popov, *Phys. Lett. A* **330**, 1 (2004); *JETP Lett.* **80**, 382 (2004); **102**, 9 (2006); S. S. Bulanov, A. M. Fedotov, and F. Pegoraro, *JETP Lett.* **80**, 734 (2004); *Phys. Rev. E* **71**, 016404 (2005).
- [15] A. R. Bell and J. G. Kirk, *Phys. Rev. Lett.* **101**, 200403 (2008); A. R. Bell, J. G. Kirk, and I. Arka, *Plasma Phys. Control Fusion* **51**, 085008 (2009).
- [16] A. M. Fedotov, N. B. Narozhny, G. Mourou, and G. Korn, *Phys. Rev. Lett.* **105**, 080402 (2010); N. V. Elkina, A. M. Fedotov, I. Yu. Kostyukov, M. V. Legkov, N. B. Narozhny, E. N. Nerush, and H. Ruhl, *Phys. Rev. ST Accel. Beams* **14**, 054401 (2011); E. N. Nerush, I. Yu. Kostyukov, A. M. Fedotov, N. B. Narozhny, N. V. Elkina, and H. Ruhl, *Phys. Rev. Lett.* **106**, 035001 (2011); B. King, N. Elkina, and H. Ruhl, *Phys. Rev. A* **87**, 042117 (2013); T. Grismayer, M. Vranic, J. L. Martins, R. A. Fonseca, and L. O. Silva, *Phys. Plasmas* **23**, 056706 (2016); S. Meuren, C. H. Keitel, and A. Di Piazza, *Phys. Rev. D* **93**, 085028 (2016).
- [17] S. S. Bulanov, T. Zh. Esirkepov, A. G. R. Thomas, J. K. Koga, and S. V. Bulanov, *Phys. Rev. Lett.* **105**, 220407 (2010).
- [18] Y. B. Zeldovich, *Sov. Phys. Usp.* **18**, 79 (1975).
- [19] J. Schwinger, L. L. De Raad, Jr., K. A. Milton, and W.-Y. Tsai, *Classical Electrodynamics* (Westview Press, Boulder, 1998).

- [20] Brown and Kibble, *Phys. Rev.* **133**, A705 (1964); C. Harvey, T. Heinzl, A. Ilderton, and M. Marklund, *Phys. Rev. Lett.* **109**, 100402 (2012).
- [21] S. J. Brodsky, H. C. Pauli, and S. S. Pinsky, *Phys. Rep.* **301**, 299 (1998); T. Heinzl, *Lect. Notes Phys.* **572**, 55 (2001).
- [22] A. Di Piazza, *Lett. Math. Phys.* **83**, 305 (2008); C. Harvey, T. Heinzl, and M. Marklund, *Phys. Rev. D* **84**, 116005 (2011); T. Heinzl, C. Harvey, A. Ilderton, M. Marklund, S. S. Bulanov, S. Rykovanov, C. B. Schroeder, E. Esarey, and W. P. Leemans, *Phys. Rev. E* **91**, 023207 (2015).
- [23] M. Vranic, J. L. Martins, J. Vieira, R. A. Fonseca, and L. O. Silva, *Phys. Rev. Lett.* **113**, 134801 (2014).
- [24] F. V. Hartemann, *High-Field Electrodynamics* (CRC Press, Boca Raton, 2002).
- [25] C. P. Ridgers, J. G. Kirk, R. Duclous, T. G. Blackburn, C. S. Brady, K. Bennett, T. D. Arber, and A. R. Bell, *J. Comput. Phys.* **260**, 273 (2014).
- [26] A. Di Piazza, K. Z. Hatsagortsyan, and C. H. Keitel, *Phys. Rev. Lett.* **105**, 220403 (2010); V. Dinu, C. Harvey, A. Ilderton, M. Marklund, and G. Torgrimsson, *Phys. Rev. Lett.* **116**, 044801 (2016).
- [27] A. Gonoskov, S. Bastrakov, E. Efimenko, A. Ilderton, M. Marklund, I. Meyerov, A. Muraviev, A. Sergeev, I. Surmin, and E. Wallin, *Phys. Rev. E* **92**, 023305 (2015).
- [28] A. M. Fedotov, arXiv:1608.02261, and references therein; V. Ritus, *Sov. Phys. JETP* **30**, 1181 (1970); N. B. Narozhny, *Phys. Rev. D* **20**, 1313 (1979); **21**, 1176 (1980).
- [29] D. Seipt and B. Kämpfer, *Phys. Rev. D* **85**, 101701 (2012); F. Mackenroth and A. Di Piazza, *Phys. Rev. Lett.* **110**, 070402 (2013); B. King, *Phys. Rev. A* **91**, 033415 (2015).
- [30] A. A. Mironov, N. B. Narozhny, and A. M. Fedotov, *Phys. Lett. A* **378**, 3254 (2014).
- [31] M. Jirka, O. Klimo, S. V. Bulanov, T. Zh. Esirkepov, E. Gelfer, S. S. Bulanov, S. Weber, and G. Korn, *Phys. Rev. E* **93**, 023207 (2016).
- [32] N. B. Narozhny, A. I. Nikisov, and V. I. Ritus, *Sov. Phys. JETP* **20**, 622 (1965).
- [33] J. Bergou and S. Varro, *J. Phys. A* **14**, 2281 (1981).